

The Influence of the Xymedon Preparation on the Regeneration of *Girardia tigrina* Planarians

Porfiriev A., Yuganova K., Vyshtakaliuk A., Zobov V., Reznik V.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Synthesis of new medicines and analysis of their effects have become increasingly important lately. Hepatoprotectors is a general name of medicinal preparations for liver treatment and recovery. The Xymedon preparation (Hydroxyethyl dimethyldihydropyrimidine), an immunostimulant, can be presumably used as a hepatoprotector. To date, the influence of this preparation has been insufficiently studied. The purpose of this paper is to investigate the effect of the pyrimidine derivative Xymedon on the regeneration of *Girardia tigrina* (Girard, 1850) planarians, one of the perspective models for analysis of the influence produced by various substances and preparations on proliferation rates of the regeneration bud (blastema). The study revealed that the Xymedon preparation is not toxic for *G. tigrina* planarians and does not cause their death even when its concentration is gradually increased in the solution. The selected working concentrations of the Xymedon preparation produced different effects on planarians during the experiment. The influence of small concentrations (0.0004 and 0.004 mg) was little different from that in the control group, i.e., the regeneration criterion was low. Higher concentrations (0.1 mg) stimulated regeneration processes, i.e., the regeneration criterion was high. The largest concentration of the Xymedon preparation (0.4 mg) led to a slight decrease of the regeneration criterion, being, however, characterized by the high regeneration coefficient.

<http://dx.doi.org/10.1007/s12668-016-0311-0>

Keywords

Hepatoprotectors, Planarians, Regeneration, Xymedon

References

- [1] Romantsov, M. G., Sukhanov, D. S., Petrov, A. Y. (2011). The use of substrates of energy metabolism in chronic liver disease for correction of metabolic disorders (experimental and clinical studies). *Fundam. Issled*, 3, 131-141.
- [2] Izmailov, S. G., Izmajlov, G. A., Aver'yanov, M. Y., Reznik, V. S. (2001). Xymedon in clinical practice (in Russian). Nizhny Novgorod.
- [3] Vyshtakaliuk, A. B., Nazarov, N. G., Porfiriev, A. G., Zueva, I. V., Minnechanova, O. A., Mayatina, O. V., et al. (2015). The influence of the xymedon preparation (hydroxyethyl dimethyldihydropyrimidine) on the rat liver recovery under toxic damage induced by carbon tetrachloride. *Doklady Biochemistry and Biophysics*, 462(1), 143-146.

- [4] Sanchez Alvarado, A., Newmark, P. A., Robb, S. M., Juste, R. (2002). The *Schmidtea mediterranea* database as a molecular resource for studying platyhelminthes, stem cells and regeneration. *Development*, 129, 5659-5665.
- [5] Tiras, K. P. (1986). Morphogenesis and ways of regeneration in planarians. *Zhurnal Obshchei Biologii*, 47(1), 103-109.
- [6] Ermakova, O. N., Ermakov, A. M., Tiras, K. P., Lednev, V. V. (2009). Effect of melatonin on regeneration of the planarian *Girardia tigrina*. *Ontogenez*, 6, 466-469.
- [7] Tiras, K. P., Petrov, O. N., Myakisheva, S. N., Aslanidi, K. B. (2015). The formation of the regeneration blastema in planarians *Dugesia (Girardia) tigrina*. *Fundam Issled*, 7(3), 493-500.
- [8] Kutsenko, S. A. (2004). *Bases of toxicology*. St. Petersburg: Foliant Publ. House.
- [9] Rombke, J., Moltmann, J.F. (1996). *Applied ecotoxicology*. CRC Lewis Publishers: Boca Raton.
- [10] Izmailov, S. G., Parshikov, V. V. (2002). Xymedon: present and future. *Nizhegorod Med Zh*, 3, 81-87.
- [11] Morawska, E., Moraczewski, J., Malczewska, M., Duma, A. (1981). Adenylate cyclase in regenerating tissues of the planarian *Dugesia lugubris* (O. Schmidt). *Hydrobiologia*, 84, 209-212.